arXiv:1204.2693v1 [math.AT] 12 Apr 2012

A nice acyclic matching on the nerve of the partition lattice

Ralf Donau[a]

[a]*Fachbereich Mathematik, Universität Bremen, Bibliothekstraße 1, 28359 Bremen, Germany*

Abstract

The author has already proven that the space $\Delta(\Pi_n)/G$ is homotopy equivalent to a wedge of spheres of dimension $n-3$ for all natural numbers $n \geq 3$ and all subgroups $G \subset S_1 \times S_{n-1}$. We wish to construct an acyclic matching on $\Delta(\Pi_n)/G$ that allows us to give a basis of its cohomology. This is also a more elementary approach to determining the number of spheres. Furthermore we give a description of the group action by an action on the spheres. We also obtain another result that we call Equivariant Patchwork Theorem.

Keywords: Discrete Morse Theory, Regular trisp, Acyclic matching, Equivariant homotopy

---

1. Introduction

Let $n \geq 3$ and let Π_n denote the poset consisting of all partitions of $[n] := \{1, \ldots, n\}$ ordered by refinement, such that the finer partition is the smaller partition. Let $\overline{\Pi}_n$ denote the poset obtained from Π_n by removing both the smallest and greatest element, which are $\{\{1\}, \ldots, \{n\}\}$ and $\{[n]\}$, respectively. We consider $\overline{\Pi}_n$ as a category, which is acyclic, and define $\Delta(\overline{\Pi}_n)$ to the nerve of the acyclic category $\overline{\Pi}_n$, which is a regular trisp, see [9, Chapter 10]. The symmetric group S_n operates on $\Delta(\overline{\Pi}_n)$ in a natural way.

It is well-known that $\Delta(\overline{\Pi}_n)$ is homotopy equivalent to a wedge of spheres of dimension $n-3$. The following two theorems are the first results concerning the topology of the quotient $\Delta(\overline{\Pi}_n)/G$, where G is a non-trivial subgroup of S_n.

Theorem 1.1 (Kozlov, [8]). *For any $n \geq 3$, the topological space $\Delta(\overline{\Pi}_n)/S_n$ is contractible.*

We set $S_1 \times S_{n-1} := \{\sigma \in S_n \mid \sigma(1) = 1\}$.

Theorem 1.2 (Donau, [2]). *Let $n \geq 3$ and $G \subset S_1 \times S_{n-1}$ be a subgroup, then the topological space $\Delta(\overline{\Pi}_n)/G$ is homotopy equivalent to a wedge of k spheres of dimension $n-3$, where k is the index of G in $S_1 \times S_{n-1}$.*

Email address: `ruelle@math.uni-bremen.de` (Ralf Donau)

This leads to a general question of determining the homotopy type of $\Delta(\overline{\Pi}_n)/G$ for an arbitrary subgroup $G \subset S_n$. One might conjecture that $\Delta(\overline{\Pi}_n)/G$ is homotopy equivalent to a wedge of spheres for any $n \geq 3$ and any subgroup $G \subset S_n$. But unfortunately this statement is not true as the following example will show.

Let $p \geq 5$ be a prime number and let C_p denote the subgroup of S_p that is generated by the cycle $(1, 2, \ldots, p)$. Then the fundamental group of $\Delta(\overline{\Pi}_p)/C_p$ is isomporphic to $\mathbb{Z}/p\mathbb{Z}$. In particular $\Delta(\overline{\Pi}_p)/C_p$ cannot be homotopy equivalent to a wedge of spheres. The proof, which uses facts about covering spaces[1], can be found in [3].

We obtain a description of the critical simplices. This result is a generalization of Lemma 4.2 in [1], where $S_1 \times S_{n-1}$ is considered.

2. Discrete Morse Theory

The definitions of regular trisps, partial matchings, acyclic matchings and foundations of Discrete Morse Theory can be found in [4, 8, 9]. The following two theorems of Discrete Morse Theory are frequently used in our proofs.

Theorem 2.1 (Patchwork Theorem). *Let $\varphi : P \longrightarrow Q$ be an order-preserving map and assume we have acyclic matchings on the subposets $\varphi^{-1}(q)$ for all $q \in Q$. Then the union of these matchings is an acyclic matching on P.*

Theorem 2.2. *Let Δ be a finite regular trisp and let M be an acyclic matching on the poset $\mathcal{F}(\Delta) \setminus \{\hat{0}\}$. Let c_i denote the number of critical i-dimensional simplices of Δ. Then Δ is homotopy equivalent to a CW complex with c_i cells of dimension i.*

The proofs of Theorems 2.1 and 2.2 as well as further facts on Discrete Morse Theory can be found in [9, Chapter 11].

Definition 2.3. *Let P be a poset and let G be a group acting on P. Let M be an acyclic matching on P. We call M an G-*equivariant acyclic matching *if $(a, b) \in M$ implies $(ga, gb) \in M$ for all $g \in G$ and $a, b \in P$.*

Remark 2.4. *Let P be a poset and let G be a finite group acting on P. Assume we have a G-equivariant acyclic matching M on P. Then M/G is an acyclic matching on P/G.*

Theorem 2.5 (Freij, [5]). *Let G be a finite group. Let Δ be a finite regular G-trisp and let M be a G-equivariant acyclic matching on the poset $\mathcal{F}(\Delta) \setminus \{\hat{0}\}$. Let c_i denote the number of critical i-dimensional simplices of Δ. Then Δ is G-homotopy equivalent to a G-CW complex where the cells correspond to the critical simplices of M and the action of G is the same as the action on Δ restricted to the critical simplices of M.*

---

[1] See [7, Chapter 1.3]

3. An Equivariant Patchwork Theorem

We wish to construct an equivariant acyclic matching on a poset by glueing together smaller equivariant acyclic matching on parts of the poset. This is similar to Theorem 2.1 with the difference that we also create copies of these matchings, see Figure 1.

Let G be a group acting on some posets P and Q. For an element $q \in Q$ we set $G_q := \{g \in G \mid gq = q\}$, known as the stabilizer subgroup of q.

Proposition 3.1. *Let $\varphi : P \longrightarrow Q$ be an order-preserving G-map and let $R \subset Q$ be a subset such that R contains exactly one representative for each orbit in Q. Assume for each $r \in R$ we have an acyclic matching M_r on $\varphi^{-1}(r)$ such that $(a, b) \in M_r$ implies $(ga, gb) \in M_r$ for all $g \in G_r$ and all $a, b \in P$. For $r \in R$, let C_r denote the set of critical elements of M_r. Then we have an G-equivariant acyclic matching on P such that*

$$\bigcup_{g \in G, r \in R} gC_r$$

is the set of critical elements.

Let $r \in R$ and assume G_r acts transitively on C_r. Then G acts transitively on

$$\bigcup_{g \in G} gC_r$$

Proof. We define acyclic matchings on the fibers of φ as follows. For each $q \in Q$ we choose $r \in R$ and $g \in G$ with $gr = q$. The map $g : \varphi^{-1}(r) \longrightarrow \varphi^{-1}(q)$, which is an isomorphism of posets, induces an acyclic matchting on $\varphi^{-1}(q)$. If we choose another $h \in G$ with $hr = q$, then we obtain the same matching. By Theorem 2.1 the union of these acyclic matchings is an acyclic matching which is G-equivariant by construction. The second statement is easy to see. $\square$

4. The main result

Let $n \geq 3$ be a fixed natural number.

Definition 4.1. *Let A be the set of all vertices of $\Delta(\overline{\Pi}_n)$ where all blocks not containing 1 are singleton. We define the following set of simplices of $\Delta(\overline{\Pi}_n)$, see Figure 2.*

$$C_n := \{\sigma \in \mathcal{F}(\Delta(\overline{\Pi}_n)) \mid V(\sigma) \subset A \text{ and } \dim \sigma = n - 3\}$$

$V(\sigma)$ denotes the set of vertices of σ and $\mathcal{F}(\Delta(\overline{\Pi}_n))$ denotes the face poset of $\Delta(\overline{\Pi}_n)$. Furthermore we set α_n to the vertex $\{\{1\}, \{2, \ldots, n\}\}$.

Remark 4.2. *The cardinality of C_n is $(n - 1)!$.*



Figure 1: An $\mathbb{Z}_2$-equivariant acyclic matching composed of $\mathbb{Z}_2$-equivariant acyclic matchings on the fibers of 0 and 1. The matching pair in the fiber of 1 is copied to the fiber of 2. $\mathbb{Z}_2$ acts on both posets by reflection across the vertical line.



Figure 2: A simplex in C_5 which has dimension 2.

Proposition 4.3. *There exists a* $(S_1 \times S_{n-1})$*-equivariant acyclic matching on* $\mathcal{F}(\Delta(\overline{\Pi}_n))$*, such that* $C_n \cup \{\alpha_n\}$ *is the set of critical simplices.*

Let V be the set of all vertices where the block containing 1 has exactly two elements and any other block is singleton. Such a vertex can be written as

$$v_k := \{\{1, k\}, \{2\}, \ldots, \widehat{\{k\}}, \ldots, \{n\}\}$$

with $k \in \{2, \ldots, n\}$. The element with the hat above is omitted.

We define a Poset $P := V \cup \{0\}$ such that 0 is the smallest element of P and the only element that is comparable with some other element. That means $x, y \in P$, $x < y$ implies $x = 0$. We define an order-preserving map $\varphi : \mathcal{F}(\Delta(\overline{\Pi}_n)) \longrightarrow P$ as follows. Let $\sigma \in \mathcal{F}(\Delta(\overline{\Pi}_n))$, then we map σ to 0 if $V(\sigma) \cap V = \emptyset$. Otherwise we map σ to the special vertex of V that belongs to σ, which is unique. Notice that $S_1 \times S_{n-1}$ acts on P in a natural way and φ is a $(S_1 \times S_{n-1})$-map. P has two orbits where one consists of one element which is 0. The other orbit may be represented by v_n.

Lemma 4.4. *There exists an* $(S_1 \times S_{n-1})$*-equivariant acyclic matching on* $\varphi^{-1}(0)$*, such that* α_n *is the only critical simplex.*

Proof. The proof of Lemma 4.4 is the same as the proof of Lemma 3.2 in [2] for the case $G = \{\mathrm{id}_{[n]}\}$. It is easy to see that the acyclic matching, that is constructed in this proof, is $(S_1 \times S_{n-1})$-equivariant. $\square$

Proof of Proposition 4.3. It is easy to see, that the statement is true for $n = 3$. Now we assume $n > 3$ and proceed by induction.

We define acyclic matchings on $\varphi^{-1}(0)$ and $\varphi^{-1}(v_n)$ as follows. By Lemma 4.4 there exists an $(S_1 \times S_{n-1})$-equivariant acyclic matching on $\varphi^{-1}(0)$, where α_n is the only critical simplex.

We define a map

$$\psi : \mathcal{F}(\Delta(\overline{\Pi}_{n-1})) \longrightarrow \varphi^{-1}(v_n) \setminus \{v_n\}$$

as follows. We add n to the block that contains 1 in each partition and append v_n to the bottom of the chain, see Figure 3. The map ψ is an isomorphism of posets. A more general definition of ψ as well as a detailed description of its inverse can be found in the proof of Lemma 4.1 in [2].



Figure 3: Example for the map ψ, where $n = 5$.

Via ψ we get an acyclic matching M on $\varphi^{-1}(v_n)$, where the set of critical simplices consists of the simplices in $\psi[C_{n-1}]$, one critical simplex s_n consisting of the two vertices v_n and $\{\{1, n\}, \{2, \ldots, (n-1)\}\}$, which has dimension 1. Additionally we have the critical simplex that has only the vertex v_n. Finally we match v_n with s_n.

We have to show that $\sigma(v_n) = v_n$ and $(a, b) \in M$ implies $(ga, gb) \in M$ for all $\sigma \in S_1 \times S_{n-1}$ and all $a, b \in \varphi^{-1}(v_n)$. Let $\sigma \in S_1 \times S_{n-1}$. $\sigma(v_n) = v_n$ implies $\sigma(1) = 1$ and $\sigma(n) = n$. We define a $\widetilde{\sigma} \in S_1 \times S_{n-2}$ by setting $\widetilde{\sigma}(x) := \sigma(x)$ for $1 \leq x \leq n-1$. Notice $\sigma\psi = \psi\widetilde{\sigma}$ which implies $\psi^{-1}\sigma = \widetilde{\sigma}\psi^{-1}$. Let $(a, b) \in M$. Clearly we have $(v_n, s_n) = (\sigma v_n, \sigma s_n)$, hence we assume $a \neq v_n$ and $b \neq s_n$. By the induction hypothesis, we have an acyclic matching $\widetilde{M}$ on $\mathcal{F}(\Delta(\overline{\Pi}_{n-1}))$ which is $(S_1 \times S_{n-2})$-equivariant. By the construction of M we have $(\psi(a)^{-1}, \psi(b)^{-1}) \in \widetilde{M}$. This implies $(\widetilde{\sigma}\psi(a)^{-1}, \widetilde{\sigma}\psi(b)^{-1}) \in \widetilde{M}$, hence $(\sigma a, \sigma b) \in M$.

By Proposition 3.1 there exists an $(S_1 \times S_{n-1})$-equivariant acyclic matching on $\mathcal{F}(\Delta(\overline{\Pi}_n))$ such that

$$\left(\bigcup_{g \in S_1 \times S_{n-1}} g\psi[C_{n-1}] \right) \cup \{\alpha_n\}$$

is the set of critical elements. It is easy to see that this set equals $C_n \cup \{\alpha_n\}$. $\square$

5. Applications

Let $n \geq 3$ be a fixed natural number.

Corollary 5.1. *The topological space $\Delta(\overline{\Pi}_n)$ is $(S_1 \times S_{n-1})$-homotopy equivalent to a wedge of $(n-1)!$ spheres of dimension $n-3$. The spheres are indexed with the simplices in C_n, which induces an action of $S_1 \times S_{n-1}$ on the $(n-1)!$ spheres.*

Proof. Apply Theorem 2.5. $\square$

Lemma 5.2. *$S_1 \times S_{n-1}$ acts freely and transitively on C_n.*

Proof. Since $C_n = \bigcup_{g \in S_1 \times S_{n-1}} g\psi[C_{n-1}]$, the action is transitive by the second statement of Proposition 3.1.

By Remark 4.2 the cardinality of C_n is $(n-1)!$ which equals the cardinality of $S_1 \times S_{n-1}$. Hence the action is free. $\square$

Let $G \subset S_1 \times S_{n-1}$ be an arbitrary subgroup.

Corollary 5.3. *There exists an acyclic matching on $\mathcal{F}(\Delta(\overline{\Pi}_n))/G$, such that the set of critical simplices consists of the simplices in C_n/G and α_n/G.*

Proof. Apply Remark 2.4. $\square$

Remark 5.4. *The cardinality of C_n/G is the index of G in $S_1 \times S_{n-1}$.*

Now Theorem 1.2 follows as a corollary:

Corollary 5.5. *The topological space $\Delta(\overline{\Pi}_n)/G$ is homotopy equivalent to a wedge of spheres of dimension $n-3$. The number of spheres is the index of G in $S_1 \times S_{n-1}$. $\Delta(\overline{\Pi}_n)^G$ is contractible for $G \neq \{id_{[n]}\}$.*

Example 5.6. *Assume $G = S_1 \times S_{n-1}$. The vertices of $\Delta(\overline{\Pi}_n)/S_1 \times S_{n-1}$ can be indexed with number partitions, which we may write as $v_0 \oplus v_1 + \cdots + v_r$, of n that distinguish the first number, i.e. $\oplus$ is non-commutative. The number on the left side of $\oplus$, that is v_0, corresponds to the block that contains 1. There exists an acyclic matching on the poset $\mathcal{F}(\Delta(\overline{\Pi}_n)/S_1 \times S_{n-1})$, where the set of critical simplices consists of the vertex $1 \oplus (n-1)$ and the unique simplex σ with the vertices $v_0 \oplus 1^{n-v_0}$ with $v_0 = 2, \ldots, n-1$ which has dimension $n-3$.*

A slightly different proof of this result, as well as a detailed description of $\Delta(\overline{\Pi}_n)/S_1 \times S_{n-1}$, can be found in [1].

Acknowledgments

The author would like to thank Dmitry N. Kozlov for this interesting problem, Ragnar Freij and Giacomo d'Antonio for the helpful discussions.

References

[1] R. Donau, *On a quotient topology of the partition lattice with forbidden block sizes*, Topology and its Applications **159** (8) (2012), pp. 2052-2057.

[2] R. Donau, *Quotients of the order complex $\Delta(\overline{\Pi}_n)$ by subgroups of the Young subgroup $S_1 \times S_{n-1}$*, Topology and its Applications **157** (16) (2010), pp. 2476-2479.

[3] R. Donau, *Quotients of the order complex $\Delta(\overline{\Pi}_n)$ that are not homotopy equivalent to wedges of spheres*, `arXiv:1202.4368v1 [math.AT]` (2012).

[4] R. Forman, *Morse theory for cell complexes*, Adv. Math. **134** (1) (1998), pp. 90-145.

[5] R. Freij, *Equivariant discrete Morse theory*, Discrete Mathematics **309** (12) (2009), pp. 3821-3829.

[6] J.P. May, *Equivariant Homotopy and Cohomology Theory*, American Mathematical Society, 1996.

[7] A. Hatcher, *Algebraic Topology*, Cambridge University Press, 2008.

[8] D.N. Kozlov, *Closure maps on regular trisps*, Topology and its Applications **156** (15) (2009), pp. 2491-2495.

[9] D.N. Kozlov, *Combinatorial Algebraic Topology*, Algorithms and Computation in Mathematics **21**, Springer-Verlag Berlin Heidelberg, 2008.

[10] A. Robinson, S. Whitehouse, *The tree representation of Σ_{n+1}*, Journal of Pure and Applied Algebra **111** (1996), pp. 245-253.